

# SOME INFINITE SERIES RELATED TO FEYNMAN DIAGRAMS

ODD MAGNE OGREID AND PER OSLAND

ABSTRACT. Results are presented for some infinite series appearing in Feynman diagram calculations, many of which are similar to the Euler series. These include both one-, two- and three-dimensional series. The sums of these series can be evaluated with the help of various integral representations for hypergeometric functions, and expressed in terms of  $\zeta(2)$ ,  $\zeta(3)$ , the Catalan constant  $G$  and  $\text{Cl}_2(\pi/3)$  where  $\text{Cl}_2(\theta)$  is Clausen's function.

## 1. INTRODUCTION

High-precision calculations in Quantum Field Theory require the evaluation of Feynman amplitudes, which represent quantum corrections in a perturbative expansion. This expansion is governed by a small parameter, the fine-structure constant, and related to the number of loops in the associated Feynman diagrams. These are graphical devices to aid in the book-keeping.

The evaluation of such Feynman amplitudes can technically be rather difficult, and one must often resort to approximate methods, valid in certain kinematical limits. For the case of electron-positron scattering, the full second-order corrections are not yet known. However, in the limit of high energies and small scattering angles, the problem simplifies: the relevant part of the so-called virtual amplitude can be expanded in powers of large logarithms. The coefficients and arguments of these logarithms can be extracted by the use of suitable Mellin transforms. We shall here review some of the techniques involved, and present some of the mathematical results of such calculations.

## 2. PHYSICS PROBLEM

The mathematical results we will discuss, derive from the “two-loop” Feynman diagram depicted in Fig. 1. The solid lines represent an electron and a positron scattering via the exchange of photons (wiggly lines). The momenta  $p_1$ ,  $p_2$ ,  $p'_1$  and  $p'_2$  only enter through the combinations

$$(2.1) \quad s = (p_1 + p_2)^2, \quad t = (p_1 - p'_1)^2 = (p_2 - p'_2)^2, \quad m^2 = p_i^2 = p_i'^2$$

where  $s$  and  $-t$  represent squares of the over-all energy and momentum transfer, respectively. Furthermore,  $m$  is the electron mass. The corresponding Feynman

---

1991 *Mathematics Subject Classification*. Primary 40A25, 40B05; Secondary 11M99, 33B15, 33C20, 33E20, 81Q30.

*Key words and phrases*. Euler series, hypergeometric series, Riemann zeta function, psi function, polylogarithms, Clausen's function.

This research has been supported by the Research Council of Norway.




FIGURE 1. Double-box diagram

amplitude is proportional to the integral

$$(2.2) \quad M = \int \frac{d^7\alpha \delta(1 - \sum \alpha_k) \Lambda^{-3}(\alpha)}{[D_s(\alpha)s + D_t(\alpha)t + D_m(\alpha)m^2 + D_\lambda(\alpha)\lambda^2]^3}$$

where a fictitious photon mass,  $\lambda$ , has been introduced in order to make the integrand well-defined for all  $\alpha_j$ . The  $D_i(\alpha)$  and  $\Lambda(\alpha)$  are homogeneous functions of seven  $\alpha_j$ , one associated with each internal line in the diagram.

Asymptotically, for  $s \gg |t| \gg m^2 \gg \lambda^2$  (high energies, small angles), the integral  $M$  has the structure

$$(2.3) \quad M \simeq A_4 \log^4(\dots) + A_3 \log^3(\dots) + \dots$$

where the arguments of the logarithms are given by the kinematical variables  $s$ ,  $t$ ,  $m^2$  and  $\lambda^2$ .

### 3. MATHEMATICAL APPROACH

**3.1. Mellin Transform.** In order to extract, from the integral (2.2), the asymptotic behaviour given by Eq. (2.3), we perform a Mellin transform. If the function  $f(x)$  behaves “almost” like  $x^{-l}$  at large  $x$ , then we define the Mellin transform as

$$(3.1) \quad J(z) = \int_0^\infty x^{l-z} f(x) dx$$

The fact that  $x^{l-z} f(x)$  is only marginally integrable, is represented by poles in  $J(z)$ , as  $z \rightarrow 0$ . The inverse transform is given by:

$$(3.2) \quad f(x) = \frac{1}{x^{l+1}} \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} x^z J(z) dz, \quad c > 0$$

where the contour of integration is as depicted in Fig. 2.

It is instructive to consider the simple example

$$(3.3) \quad f(x) = \theta(x-1) \frac{1}{x^{l+1}} \log^k x,$$

for which the Mellin transform gives

$$(3.4) \quad J(\zeta) = \frac{k!}{\zeta^{k+1}}$$

Thus, an expansion of the Mellin transform in inverse powers of  $\zeta$  is equivalent to an expansion in powers of logarithms.




FIGURE 2. Contour used for the Inverse Mellin transform

**3.2. Factorization Formula.** In order to facilitate the integration over the  $\alpha_j$  in (2.2), we invoke the “factorization formula” [1, 12]. For three terms, it can be written as:

$$(3.5) \quad \begin{aligned} & (A_1 + A_2 + A_3)^{-p} \\ &= \int_{c_1} \frac{dz_1}{2\pi i} \int_{c_2} \frac{dz_2}{2\pi i} \frac{\Gamma(z_1)\Gamma(z_2)\Gamma(p-z_1-z_2)}{\Gamma(p)} A_1^{-z_1} A_2^{-z_2} A_3^{-p+z_1+z_2} \\ & \quad \text{Im } A_i > 0, \quad c_j > 0, \quad p - c_1 - c_2 > 0 \end{aligned}$$

where the integration contours labelled  $c_j$  run from  $c_j - i\infty$  to  $c_j + i\infty$ . To prove the factorization formula, one will need two preliminary results. The first one is

$$(3.6) \quad u^{-p} = \frac{e^{-i\pi p/2}}{\Gamma(p)} \int_0^\infty t^{p-1} e^{itu} dt, \quad \text{Re } p > 0, \quad \text{Im } u > 0.$$

*Proof:* First, change variables by putting  $x = -itu$ . Using the integral representation of the Gamma function, we find,

$$\frac{e^{-i\pi p/2}}{\Gamma(p)} \int_0^{-iu\infty} (-iu)^{-p} e^{-x} x^{p-1} dx = \frac{e^{-i\pi p/2}}{\Gamma(p)} e^{i\pi p/2} u^{-p} \Gamma(p) = u^{-p}$$

which is valid if  $u$  is purely imaginary with  $u/i > 0$  and  $\text{Re } p > 0$ . This result can be extended to any  $u$  with  $\text{Im } u > 0$  by analytic continuation. The second result needed is

$$(3.7) \quad e^{iT} = \frac{1}{2\pi i} \int_c \Gamma(z) e^{i\pi z/2} T^{-z} dz, \quad c > 0, \quad \text{Im } T > 0.$$

*Proof:* Consider the Mellin transform of  $e^{iTx}$  [cf. Eq. (3.6)],

$$I' = \int_0^\infty x^{z-1} e^{iTx} dx = \Gamma(z) e^{i\pi z/2} T^{-z},$$

which is valid for  $\text{Re } z > 0$  and  $\text{Im } T > 0$ . By taking the inverse Mellin transform of this expression, one should get  $e^{iTx}$ ,

$$e^{iTx} = \frac{1}{2\pi i} \int_{c'} x^{-z} \Gamma(z) e^{i\pi z/2} T^{-z} dz.$$

By putting  $x = 1$  in this expression, the desired result is obtained. One is now capable of proving the factorization formula.

*Proof of the factorization formula:*

$$\begin{aligned}
(A_1 + A_2 + A_3)^{-p} &= \frac{e^{-i\pi p/2}}{\Gamma(p)} \int_0^\infty dt t^{p-1} e^{it(A_1+A_2+A_3)} \\
&= \frac{e^{-i\pi p/2}}{\Gamma(p)} \int_0^\infty dt t^{p-1} e^{itA_1} e^{itA_2} e^{itA_3} \\
&= \frac{e^{-i\pi p/2}}{\Gamma(p)} \int_0^\infty dt t^{p-1} \int_{c_1} \frac{dz_1}{2\pi i} \int_{c_2} \frac{dz_2}{2\pi i} \Gamma(z_1) \Gamma(z_2) \\
&\quad \times e^{i\pi z_1/2} e^{i\pi z_2/2} (tA_1)^{-z_1} (tA_2)^{-z_2} e^{itA_3} \\
&= \frac{e^{-i\pi p/2}}{\Gamma(p)} \int_{c_1} \frac{dz_1}{2\pi i} \int_{c_2} \frac{dz_2}{2\pi i} \Gamma(z_1) \Gamma(z_2) \\
&\quad \times e^{i\pi z_1/2} e^{i\pi z_2/2} A_1^{-z_1} A_2^{-z_2} \int_0^\infty dt t^{p-1-z_1-z_2} e^{itA_3} \\
&= \frac{e^{-i\pi p/2}}{\Gamma(p)} \int_{c_1} \frac{dz_1}{2\pi i} \int_{c_2} \frac{dz_2}{2\pi i} \Gamma(z_1) \Gamma(z_2) e^{i\pi z_1/2} e^{i\pi z_2/2} \\
&\quad \times A_1^{-z_1} A_2^{-z_2} \Gamma(p - z_1 - z_2) e^{i\pi(p-z_1-z_2)/2} A_3^{-p+z_1+z_2},
\end{aligned}$$

from which the factorization formula follows. The formula is valid for  $\text{Im } A_i > 0$ ,  $c_i > 0$  and  $p - c_1 - c_2 > 0$ . It is trivially generalized to  $(A_1 + A_2 + \dots + A_n)^{-p}$ .

**3.3. Emergence of Sums.** Performing a Mellin transform in  $s$ , with  $|t|=1$ ,  $m^2 = s^{\eta_m}$ ,  $\lambda^2 = s^{\eta_\lambda}$ , and  $\eta_\lambda < \eta_m < 0$ , one is led to integrals over  $z_i$  of the kind given by Eq. (3.5). These integrals are most conveniently performed by the use of Cauchy's theorem, closing the contour in the left or right half-plane. Expanding these summands in powers of the Mellin transform variable, one is led to sums involving, as residues, the  $\Gamma$  function and its derivatives,

$$\psi(z) = (d/dz) \log \Gamma(z) = \Gamma'(z)/\Gamma(z)$$

and

$$\psi'(z) = (d/dz) \psi(z).$$

The combination  $\gamma + \psi(n)$  frequently appears in the summand, and several series of this type are evaluated using the following integral representation,

$$\gamma + \psi(z) = \int_0^1 dt \frac{1-t^{z-1}}{1-t}.$$

Here,  $\gamma = 0.577\,216\dots$  is Euler's constant.

**3.4. Special Functions.** Our evaluation of the sums makes use of several special functions. We shall list a few of them along with some of their properties here:

*Riemann's zeta function:*

The sums of several series can be expressed in terms of the Riemann  $\zeta$  function:

$$\zeta(z) = \sum_{k=1}^{\infty} \frac{1}{k^z}$$

with

$$\sum_{k=1}^{\infty} \frac{1}{k^2} = \zeta(2) = \frac{\pi^2}{6} = 1.644\,934\dots,$$

$$\sum_{k=1}^{\infty} \frac{1}{k^3} = \zeta(3) = 1.202\,057\dots$$

*Polylogarithms:*

Furthermore, we need Euler's polylogarithms:

$$\text{Li}_n(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^n}, \quad |z| < 1, \quad n = 0, 1, 2, \dots,$$

These functions also have integral representations:

$$(3.8) \quad \text{Li}_n(z) = \frac{(-1)^{n-1}}{(n-2)!} \int_0^1 dt \frac{\log^{n-2}(t) \log(1-tz)}{t}, \quad n = 2, 3, 4, \dots$$

$$(3.9) \quad = \int_0^z dt \frac{\text{Li}_{n-1}(t)}{t}, \quad n = 1, 2, 3, \dots,$$

with  $\text{Li}_n(1) = \zeta(n)$ ,  $n = 2, 3, 4, \dots$ ,  $\text{Li}_n(-1) = (1/2^{n-1} - 1)\zeta(n)$ ,  $n = 2, 3, 4, \dots$ .

*Clausen's function:*

On the unit circle, the imaginary part of the dilogarithm is Clausen's function:

$$\text{Cl}_2(\theta) = \text{Im}[\text{Li}_2(e^{i\theta})] = \sum_{k=1}^{\infty} \frac{\sin k\theta}{k^2}.$$

Clausen's function has its maximum value for  $\theta = \pi/3$ ,  $\text{Cl}_2(\pi/3) = 1.014\,942\dots$ .

Furthermore,  $\text{Cl}_2(\pi/2) = G = 0.915\,966\dots$ , where  $G$  is Catalan's constant.

**3.5. A Theorem For Summing Series.** Many of the results given here were obtained using "Theorem 1" of [10]:

$$(3.10) \quad \sum_{n=1}^{\infty} \frac{1}{n^2} [\gamma + \psi(1+kn)] = \left( \frac{k^2}{2} + \frac{3}{2k} \right) \zeta(3) + \pi \sum_{j=1}^{k-1} j \text{Cl}_2 \left( \frac{2\pi j}{k} \right)$$

$$(3.11) \quad \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} [\gamma + \psi(1+kn)] = \left( \frac{k^2}{2} - \frac{9}{8k} \right) \zeta(3) + \pi \sum_{j=1}^{k-1} j \text{Cl}_2 \left( \frac{2\pi j}{k} + \frac{\pi}{k} \right)$$

for  $k = 1, 2, 3, \dots$ . The sums over  $j$  are understood to be zero when  $k = 1$ .

To prove the theorem, we use properties of Nielsen's generalized polylogarithms [6]. For a detailed proof of the theorem, we refer to [10].

#### 4. ONE-DIMENSIONAL SERIES

We present a collection of one-dimensional series and their sums. For some of the series we also present the methods used to sum them. For the rest of the series,

the proofs can be found in [10, 11].

$$(4.1) \quad \sum_{n=1}^{\infty} \frac{1}{n^2} [\gamma + \psi(1+n)] = 2\zeta(3)$$

$$(4.2) \quad \sum_{n=1}^{\infty} \frac{1}{n^2} [\gamma + \psi(1+2n)] = \frac{11}{4}\zeta(3)$$

$$(4.3) \quad \sum_{n=1}^{\infty} \frac{1}{n^2} [\gamma + \psi(1+3n)] = 5\zeta(3) - \frac{2\pi}{3}\text{Cl}_2\left(\frac{\pi}{3}\right)$$

$$(4.4) \quad \sum_{n=1}^{\infty} \frac{1}{n^2} [\gamma + \psi(1+4n)] = \frac{67}{8}\zeta(3) - 2\pi G$$

$$(4.5) \quad \sum_{n=1}^{\infty} \frac{1}{n^2} [\gamma + \psi(1+6n)] = \frac{73}{4}\zeta(3) - \frac{16\pi}{3}\text{Cl}_2\left(\frac{\pi}{3}\right)$$

Some of these series were known already by Euler! Alternating series give similar results:

$$(4.6) \quad \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} [\gamma + \psi(1+n)] = -\frac{5}{8}\zeta(3)$$

$$(4.7) \quad \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} [\gamma + \psi(1+2n)] = \frac{23}{16}\zeta(3) - \pi G$$

$$(4.8) \quad \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} [\gamma + \psi(1+3n)] = \frac{33}{8}\zeta(3) - 2\pi\text{Cl}_2\left(\frac{\pi}{3}\right)$$

The results presented so far, all follow from the theorem or the immediate corollary thereof.

Some series encountered had to be summed by other means:

$$(4.9) \quad \sum_{n=1}^{\infty} \frac{1}{n(n+1)} [\gamma + \psi(1+n)] = \zeta(2)$$

$$(4.10) \quad \sum_{n=1}^{\infty} \frac{1}{n^2(n+1)} [\gamma + \psi(1+n)] = 2\zeta(3) - \zeta(2)$$

$$(4.11) \quad \sum_{n=1}^{\infty} \frac{1}{n(n+1)^2} [\gamma + \psi(1+n)] = -\zeta(3) + \zeta(2)$$

Related sums, where the argument of the psi function is shifted by a small integer, are easily obtained using the recursion relation  $\psi(1+n) = \psi(n) + 1/n$ .

*Proof of (4.9):* We apply the integral representation of the psi function before

summing over  $n$ :

$$\begin{aligned}
\sum_{n=1}^{\infty} \frac{1}{n(n+1)} [\gamma + \psi(1+n)] &= \sum_{n=1}^{\infty} \frac{1}{n(n+1)} \int_0^1 dt \frac{1-t^n}{1-t} \\
&= \int_0^1 dt \frac{1}{1-t} \left[ 1 - \frac{t}{2} {}_2F_1(1, 1; 3; t) \right] \\
&= \int_0^1 dt \frac{1}{1-t} \left\{ 1 - \frac{1}{t} [t + (1-t) \log(1-t)] \right\} \\
&= - \int_0^1 dt \frac{\log(1-t)}{t} = \zeta(2)
\end{aligned}$$

Here, we used (7.3.2.150) of Prudnikov [14] to rewrite  ${}_2F_1$ . The resulting integral is just the definition of  $\text{Li}_2(1)$  which is well known.

Series containing the derivative of the psi function were also encountered:

$$(4.12) \quad \sum_{n=1}^{\infty} \frac{1}{n} \psi'(n) = 2\zeta(3)$$

$$(4.13) \quad \sum_{n=1}^{\infty} \frac{1}{n} \psi'(1+n) = \zeta(3)$$

$$(4.14) \quad \sum_{n=1}^{\infty} \frac{1}{n(n+1)} \psi'(n) = 1$$

$$(4.15) \quad \sum_{n=1}^{\infty} \frac{1}{n(n+1)} \psi'(1+n) = -\zeta(3) + \zeta(2)$$

*Proof of Series (4.13):* We start by using the integral representation of the trigamma function:

$$\begin{aligned}
\sum_{n=1}^{\infty} \frac{1}{n} \psi'(1+n) &= - \sum_{n=1}^{\infty} \frac{1}{n} \int_0^1 dt \frac{t^n}{1-t} \log t \\
&= \int_0^1 \frac{dt}{1-t} \log t \log(1-t) \\
&= \int_0^1 \frac{dt}{t} \log t \log(1-t) = \zeta(3)
\end{aligned}$$

The resulting integral is just the integral representation of  $\text{Li}_3(1)$  which equals  $\zeta(3)$ .

Also some series bilinear in  $\psi(z)$  can be summed by similar methods:

$$(4.16) \quad \sum_{n=1}^{\infty} \frac{1}{n(n+1)} [\gamma + \psi(n)]^2 = \zeta(2) + 1$$

$$(4.17) \quad \sum_{n=1}^{\infty} \frac{1}{n(n+1)} [\gamma + \psi(n)][\gamma + \psi(1+n)] = \zeta(3) + \zeta(2)$$

$$(4.18) \quad \sum_{n=1}^{\infty} \frac{1}{n(n+1)} [\gamma + \psi(n)][\gamma + \psi(2+n)] = 3$$

$$(4.19) \quad \sum_{n=1}^{\infty} \frac{1}{n(n+1)} [\gamma + \psi(1+n)]^2 = 3\zeta(3)$$

$$(4.20) \quad \sum_{n=1}^{\infty} \frac{1}{n(n+1)} [\gamma + \psi(1+n)][\gamma + \psi(2+n)] = 2\zeta(3) + \zeta(2)$$

$$(4.21) \quad \sum_{n=1}^{\infty} \frac{1}{n(n+1)} [\gamma + \psi(2+n)]^2 = \zeta(2) + 3$$

These series are summed by using the integral representation of each of the factors  $\gamma + \psi(1+n)$  before summing over  $n$ . The resulting two-dimensional integral yields the desired result.

Some one-dimensional series containing  $\Gamma$ -functions also appear. Consider the following series:

$$(4.22) \quad \sum_{n=1}^{\infty} \frac{1}{n^2} \frac{[\Gamma(n)]^2}{\Gamma(2n)} = -\frac{8}{3}\zeta(3) + \frac{4\pi}{3}\text{Cl}_2\left(\frac{\pi}{3}\right)$$

$$(4.23) \quad \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} \frac{[\Gamma(n)]^2}{\Gamma(2n)} = -\frac{4}{5}\zeta(3)$$

*Proof of (4.22):*

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{1}{n^2} \frac{[\Gamma(n)]^2}{\Gamma(2n)} &= 2 \sum_{n=1}^{\infty} \frac{1}{n^2} \frac{\Gamma\left(\frac{1}{2}\right) \Gamma(n)}{\Gamma\left(\frac{1}{2}+n\right)} \left(\frac{1}{4}\right)^n \\ &= {}_4F_3\left(1, 1, 1, 1; \frac{3}{2}, 2, 2; \frac{1}{4}\right) \\ &= \int_0^1 dt {}_3F_2\left(1, 1, 1; \frac{3}{2}, 2; \frac{1}{4}t\right) \end{aligned}$$

Next, we make use of (7.4.2.353) of [14] and the substitution  $\sqrt{t}/2 = \sin \frac{u}{2}$  to get

$$4 \int_0^1 \frac{dt}{t} \arcsin^2 \frac{\sqrt{t}}{2} = 2 \int_0^{\frac{\pi}{3}} \frac{du}{2 \tan \frac{u}{2}} u^2.$$

Integration by parts yields:

$$\begin{aligned} &2 u^2 \log\left(2 \sin \frac{u}{2}\right) \Big|_0^{\frac{\pi}{3}} - 4 \int_0^{\frac{\pi}{3}} du u \log\left(2 \sin \frac{u}{2}\right) \\ &= -4 \int_0^{\frac{\pi}{3}} du u \log\left(2 \sin \frac{u}{2}\right) = -\frac{8}{3}\zeta(3) + \frac{4\pi}{3}\text{Cl}_2\left(\frac{\pi}{3}\right) \end{aligned}$$

after using (6.46) and correcting the misprint in (6.52) of [8].

## 5. TWO-DIMENSIONAL SERIES

In the evaluation of Feynman diagrams by the above outlined methods, also two- and three-dimensional series are required. For the purpose of summing certain two-dimensional series, the following results are very useful:

$$(5.1) \quad \sum_{n=1}^{\infty} \frac{1}{n+a} \frac{1}{n+b} = \frac{1}{b-a} [\psi(1+b) - \psi(1+a)], \quad a \neq b$$



and

$$(5.2) \quad \sum_{n=1}^{\infty} \frac{\Gamma(n+k)}{\Gamma(1+n+2k)} = \frac{\Gamma(k)}{\Gamma(1+2k)}$$

Some selected two-dimensional series that do not involve the psi function are:

$$(5.3) \quad \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \frac{1}{nk(n+k)} = 2\zeta(3)$$

$$(5.4) \quad \sum_{n=0}^{\infty} \sum_{k=1}^{\infty} \frac{1}{k(n+k)(1+n+k)} = \zeta(2)$$

$$(5.5) \quad \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \frac{1}{k} \frac{\Gamma(n)\Gamma(k)}{\Gamma(1+n+k)} = \zeta(3)$$

$$(5.6) \quad \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \frac{1}{k!} \frac{\Gamma(2k)\Gamma(n+k)}{\Gamma(1+n+2k)} = \frac{1}{2}\zeta(2)$$

A few typical series involving the psi function are:

$$(5.7) \quad \sum_{n=0}^{\infty} \sum_{k=1}^{\infty} \frac{\gamma + \psi(1+k)}{k(n+k)(1+n+k)} = 2\zeta(3)$$

$$(5.8) \quad \sum_{n=0}^{\infty} \sum_{k=1}^{\infty} \frac{\gamma + \psi(1+n)}{k(n+k)(1+n+k)} = 2\zeta(3)$$

$$(5.9) \quad \sum_{n=0}^{\infty} \sum_{k=1}^{\infty} \frac{\gamma + \psi(1+n+k)}{k(n+k)(1+n+k)} = 3\zeta(3)$$

$$(5.10) \quad \sum_{n=0}^{\infty} \sum_{k=1}^{\infty} \frac{\gamma + \psi(1+n+2k)}{k(n+k)(1+n+k)} = \frac{7}{2}\zeta(3)$$

## 6. SUMMARY

We have seen that Quantum Field Theory relates to rather interesting mathematics. In physical problems of central interest (see, e.g., [4, 5, 9, 13, 15]), series related to the Euler series emerge. They have been summed in terms of  $\zeta(2)$ ,  $\zeta(3)$  and closely related irrational constants. A computerized treatment of such sums is also available [16].

It is interesting to note that the transcendental basis that is required for the considered types of sums is rather restricted. In particular, there are no  $\zeta(4)$ ,  $\zeta(5)$ , etc. The restrictions of such transcendental bases have received a lot of attention recently, in particular by Broadhurst and Kreimer [2, 7] (see also [3]), who relate these restrictions to properties of Hopf algebras.

## REFERENCES

- [1] J.D. Bjorken and T.T. Wu, *Phys. Rev.* **130** (1963) 2566
- [2] D.J. Broadhurst and D. Kreimer, Association of multiple zeta values with positive knots via Feynman diagrams up to 9 loops, *Phys. Lett.* **B393** (1997) 403; J.M. Borwein and D.J. Broadhurst, Determinations of rational Dedekind-zeta invariants of hyperbolic manifolds and Feynman knots and links, preprint, OUT-4102-76 (1998), hep-th/9811173.
- [3] S. Groote, J.G. Körner and A.A. Pivovarov, Transcendental numbers and the topology of three-loop bubbles, *Phys. Rev.* **D60** (1999) 061701

- [4] S. Groote and A.A. Pivovarov, Threshold expansion of Feynman diagrams within a configuration space technique, *Nucl. Phys.* **B580** (2000) 459
- [5] M.Yu. Kalmykov and O. Veretin, Single scale diagrams and multiple binomial sums, *Phys. Lett.* **B483** (2000) 315;
- [6] K. S. Kölbig, J. A. Mignaco, and E. Remiddi, On Nielsen's generalized polylogarithms and their numerical calculation, *BIT* **10** (1970) 38
- [7] see, for example: A. Connes and D. Kreimer, Hopf algebras, renormalization and noncommutative geometry, *Commun. Math. Phys.* **199** (1998) 203; D. Kreimer, and R. Delbourgo, Using the Hopf algebra structure of QFT in calculations, *Phys. Rev.* **D60** (1999) 105025
- [8] L. Lewin, Polylogarithms and associated functions, North-Holland, New York, 1981.
- [9] K. Melnikov and T. van Ritbergen, The three loop on shell renormalization of QCD and QED, SLAC-PUB-8450, hep-ph/0005131
- [10] O. M. OGREID and P. OSLAND, Summing one- and two-dimensional series related to the Euler series, *J. Comput. Appl. Math.* **98** (1998) 245
- [11] O. M. OGREID and P. OSLAND, More Series related to the Euler Series, *J. Comput. Appl. Math.*, in print; hep-th/9904206
- [12] P. OSLAND and T.T. WU, *Nucl. Phys.* **B288** (1987) 77
- [13] A.C. Petkou and G. Siopsis, Lee-Yang edge singularity in the three-dimensional Gross-Neveu model at finite temperature, preprint KL-TH-00-05, hep-th/0008198
- [14] A.P. Prudnikov, Yu.A. Brychkov, O.I. Marichev, *Integrals and Series*, vol. 3, Gordon and Breach, New York, 1990.
- [15] T. van Ritbergen, R.G. Stuart, On the precise determination of the Fermi coupling constant from the muon lifetime, *Nucl. Phys.* **B564** (2000) 343
- [16] J. Vermaseren, *Int. Journal of Modern Phys.* **A 14** (1999) 2037.

DEPARTMENT OF PHYSICS, UNIVERSITY OF BERGEN, ALLÉGT. 55, N-5007 BERGEN, NORWAY  
*E-mail address:* Odd.Ogreid@fi.uib.no  
*URL:* <http://www.fi.uib.no/~ogreid/>

DEPARTMENT OF PHYSICS, UNIVERSITY OF BERGEN, ALLÉGT. 55, N-5007 BERGEN, NORWAY  
*E-mail address:* Per.Osland@fi.uib.no  
*URL:* <http://www.fi.uib.no/~osland/particle.html>